

Trace the Denmark Strait overflow water in an eddy-resolving Atlantic simulation: some preliminary results

Xiaobiao Xu (COAPS/FSU),
Alan Wallcraft (NRL/SSC),
Eric Chassignet (COAPS/FSU)

Thanks: Peter Rhines (UW) and William Schmitz


May 21-23, Layered ocean modeling workshop, Ann Arbor, MI

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE MAY 2013		2. REPORT TYPE		3. DATES COVERED 00-00-2013 to 00-00-2013	
4. TITLE AND SUBTITLE Trace the Denmark Strait overflow water in an eddy-resolving Atlantic simulation: some preliminary results				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Florida State University ,Center for Ocean-Atmospheric Prediction Studies (COAPS),2000 Levy Avenue, Building A, Suite 292,Tallahassee,FL,32306-2741				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 18	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

Motivation


- Passive tracer, when tagged to a particular water mass, can be used to “effectively” track the movement/circulation of the water mass and its interaction with the ambient water.
- When the Denmark Strait overflow water (DSOW) flow into the Irminger and Labrador Seas, the observed vertical structure of horizontal velocity shows a strong bottom intensification, typically not in general circulation models. We want to use tracer to quantify the associated mixing and water mass transformation.
- Tracers (CFC, Oxygen, I^{129} , etc) have been widely used in observations associated with the DSOW, e.g., Smith et al., 2005. Can models simulate those tracer signature?

3-layer circulation in the SPNA




upper layer: Red and yellow ("modified" North Atlantic Water); middle layer: green (Labrador Sea Water); deep layer: blue (Nordic Seas Overflow Water)

Overflow structure in HYCOM vs. obs.




(a) Angmagssalik




(b) Southeast of Cape Farewell

Comparison near 53N



Moored observations from Fischer et al., 2004, 2010. HYCOM results and comparison to observations see Xu et al., 2013

Comparison near 43N




Moored observations from Clarke et al., 1998; Meinen 2000; Schott et al., 2004, 2006. Model results and comparison to observations see Xu et al., 2013

Tracer in the Atlantic Simulation


- HYbrid Coordinate Ocean Model (HYCOM, Bleck, 2002; Chassignet et al., 2003; <http://www.hycom.org>); see Xu et al., 2010, 2012, 2013 for documentation of the Atlantic model results
- Model domain extends meridionally from 28° S to 80° N, as a subset of the global prediction system developed at NRL/SSC
- 0.08° horizontal resolution (5-6 km in the SPNA); 32 σ_2 layers in vertical, with thermo-baric effect parameterized
- Initialization with zero velocity and January potential temperature and salinity from monthly ocean climatology GDEM (Carnes, 2009)
- Climatological forcing (E026): 25-year spin-up using forcing based on a monthly climatology (ERA40, Uppala et al., 2005) + submonthly wind anomalies from Navy Operational Global Atmospheric Prediction System (NOGAPS) for year 2003;
- Tracers were injected on Jan 1 of model year 16 (at the Denmark Strait sill below model layer 21 and in the Florida Strait in the top 15 layers). The simulation is integrated for 10 years.

10 years after (Florida Strait)



Integrated tracer content below model layers at the end of the simulation;


10 years after (Demark Strait)








Integrated tracer content below model layer 20 at the end of the simulation;

Vertical distribution (1 year mean)






Passive-tracer 67.88n - 65.45n mean: 23.955- 25.045 [03.2H]



Passive-tracer 53.67n - 60.51n mean: 23.955- 25.045 [03.2H]




Compared to the observed I^{129}




In the Labrador Sea

I129 from Smith et al., 2005




Both the observed I^{129} and numerical tracer spread out the lowest part of *entire* Labrador Sea.




Vertical structure of tracer flux


Denark Strait




Angmagssalik




Cape Farewell




Labrador Sea




Grand Banks






50W



Total tracer fluxes in overflow water




Full water column budget



Net flux through the boundaries


Inventory change in the box



Summary

- A numerical tracer experiment is conducted in an eddy-resolving Atlantic simulation.
- The DSOW tracer is seen descend from the Denmark Sill to the observed DSOW depth range in the Irminger and Labrador Sea, with a vertical distribution similar to the observed I^{129} .
- Preliminary analysis suggest that immediately after the DSOW spreading into the Irminger Sea, some tracer have been *lost* to water above the overflow.
- The full water column budget of the tracer flux/content is *approximately* conserved.
- **currently on going ** simplify the experiment by releasing tracer only in the DSOW, and save the model fluxes to test the conservation of tracer content.

If I saved the model volume fluxes ...



Thank you

